ELSEVIER

Environmental Pollution 152 (2008) I–VI

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 152

Number 1 2008

Regular papers

1 **Seasonal variation in the atmospheric deposition of inorganic constituents and canopy interactions in a Japanese cedar forest**
H. Sase, A. Takahashi, M. Sato, H. Kobayashi, M. Nakata, T. Totsuka

Leaf surface properties may contribute to the ion transport process of the forest canopy.

11 **Natural and pre-treatments induced variability in the chemical composition and morphology of lichens and mosses selected for active monitoring of airborne elements**
P. Adamo, R. Bargagli, S. Giordano, P. Modenesi, F. Monaci, E. Pittao, V. Spagnuolo, M. Tretiach

Water-washed and oven-dried moss shoots and lichen lobes have adequate chemical and morphological homogeneity to be used as active biomonitors of trace element atmospheric deposition.

20 **Effect of long-term changes in soil chemistry induced by road salt applications on N-transformations in roadside soils**
S.M. Green, R. Machin, M.S. Cresser

Road salting effects ammonification and nitrification in roadside soils.

32 **Assessment of the phytoextraction potential of high biomass crop plants**
J. Hernández-Allica, J.M. Becerril, C. Garbisu

The phytoextraction performance of cultivars varies significantly depending on the screening method used.

41 **Long-term nitrate increases in two oligotrophic lakes, due to the leaching of atmospherically-deposited N from moorland ranker soils**
E. Tipping, S.A. Thacker, D. Wilson, J.R. Hall

Increases in lakewater nitrate concentrations over 50 years are due to the limited ability of ranker soils to retain atmospherically-deposited nitrogen.

50 **Trace element accumulation in woody plants of the Guadiamar Valley, SW Spain: A large-scale phytomanagement case study**
M.T. Domínguez, T. Marañón, J.M. Murillo, R. Schulin, B.H. Robinson

There is a low trace element transfer from contaminated soils to the aboveground parts of afforested woody plants under a semi-arid climate.

60 **Use of the physiologically-based extraction test to assess the oral bioaccessibility of metals in vegetable plants grown in contaminated soil**
M. Intawongse, J.R. Dean

Evaluation of a physiologically-based extraction test to assess the risk to humans of consuming contaminated vegetables.

73 **Metals in particle-size fractions of the soils of five European cities**
F. Ajmone-Marsan, M. Biasioli, T. Kralj, H. Grčman, C.M. Davidson, A.S. Hursthouse, L. Madrid, S. Rodrigues

Metals in size fractions of urban soils.

82 **Differences in the growth response of three bryophyte species to nitrogen**
M. Salemaa, R. Mäkipää, J. Oksanen

Boreal bryophytes display differences in their sensitivity to nitrogen.

92 **Effects of repeated exposure to malathion on growth, food consumption, and locomotor performance of the western fence lizard (*Sceloporus occidentalis*)**
R.R. Holem, W.A. Hopkins, L.G. Talent

Repeated exposure of western fence lizards to malathion caused reduced arboreal performance and some mortality but growth, food consumption, and terrestrial performance were not affected.

99 **Influence of surfactant sorption on the removal of phenanthrene from contaminated soils**
W. Zhou, L. Zhu

The sorption of surfactant onto soils had a potential influence on the remediation time scales and surfactant addition strategies of surfactant-enhanced remediation for contaminated soils.

106 **Impacts of microbial redox conditions on the phase distribution of pyrene in soil–water systems**
H.S. Kim, J.C. Roper, F.K. Pfaender

HOC association within soil/sediment matrix can be controlled by microbially mediated redox conditions.

doi:10.1016/S0269-7491(08)00145-0

116 Is selenium affecting body condition and reproduction in boreal breeding scaup, scoters, and ring-necked ducks?
J.-M.A. DeVink, R.G. Clark, S.M. Slattery, M. Wayland

Selenium is not affecting boreal scaup, ring-necked duck, or scoter body condition or reproduction.

123 Construction and comparison of fluorescence and bioluminescence bacterial biosensors for the detection of bioavailable toluene and related compounds
Y.-F. Li, F.-Y. Li, C.-L. Ho, V.H.-C. Liao

The choice of reporter genes for toluene bacterial biosensors to determine BTEX bioavailability is case-specific.

130 Partitioning of polycyclic aromatic hydrocarbons to solid-sorbed nonionic surfactants
L. Zhu, W. Zhou

The sorption of solid-sorbed nonionic surfactant for PAHs may be attributed to PAHs partitioning to surface micelle (or admicelle) of surfactant on solid surface.

138 Chemical and biochemical tools to assess pollution exposure in cultured fish
D. Fernandes, S. Zanuy, M.J. Bebianno, C. Porte

Pollution assessment in cultured fish: chemical and biochemical tools.

147 Modeling uptake kinetics of cadmium by field-grown lettuce
W. Chen, L. Li, A.C. Chang, L. Wu, S.-I. Kwon, R. Bottoms

A kinetic model was developed to evaluate the uptake of Cd under field conditions.

153 Field and laboratory simulations of storm water pulses: Behavioural avoidance by marine epifauna
D.A. Roberts, E.L. Johnston, S. Müller, A.G.B. Poore

Brief storm water pulses trigger avoidance response in mobile epifauna due to the inundation of freshwater.

163 The ratio of clay content to total organic carbon content is a useful parameter to predict adsorption of the herbicide butachlor in soils
Z. Liu, Y. He, J. Xu, P. Huang, G. Jilani

The relative importance of organic matter and clay in butachlor adsorption in soil will depend on the ratio of clay content to total organic carbon content.

172 Influence of industrial heavy metal pollution on soil free-living nematode population
S. Pen-Mouratov, N. Shukurov, Y. Steinberger

Trophic structure and sex ratio of soil nematode population are sensitive tools for monitoring industrial pollution.

184 Phosphate-induced lead immobilization from different lead minerals in soils under varying pH conditions
X. Cao, L.Q. Ma, S.P. Singh, Q. Zhou

Phosphate effectively converts litharge (PbO), cerrusite ($PbCO_3$), and anglesite ($PbSO_4$) in a soil under different pHs to pyromorphite, thereby reducing Pb solubility in the soil.

193 The dispersal and storage of sediment-associated metals in an arid river system: The Leichhardt River, Mount Isa, Queensland, Australia
M.P. Taylor, K.A. Hudson-Edwards

Cu-Pb-Zn contamination of aquatic systems draining Mount Isa Mine, Australia.

205 Changes in persistent contaminant concentration and CYP1A1 protein expression in biopsy samples from northern bottlenose whales, *Hyperoodon ampullatus*, following the onset of nearby oil and gas development
S.K. Hooker, T.L. Metcalfe, C.D. Metcalfe, C.M. Angell, J.Y. Wilson, M.J. Moore, H. Whitehead

Whale contaminants highlight concerns from oil and gas development near a marine protected area.

217 Examining the joint toxicity of chlorpyrifos and atrazine in the aquatic species: *Lepomis macrochirus*, *Pimephales promelas* and *Chironomus tentans*
W. Tyler Mehler, L.J. Schuler, M.J. Lydy

The joint toxicity between atrazine and chlorpyrifos caused greater than additive responses in invertebrates, but the interactions in vertebrates was less pronounced.

225 Life cycle responses of the midge *Chironomus riparius* to polycyclic aromatic compound exposure
M.L. Paumen, E. Borgman, M.H.S. Kraak, C.A.M. van Gestel, W. Admiraal

Emergence of Chironomus riparius is a sensitive endpoint to detect life cycle effects of PACs.

233 Induced cytochrome P450 1A activity in cichlid fishes from Guandu River and Jacarepaguá Lake, Rio de Janeiro, Brazil
T.E.M. Parente, A.C.A.X. De-Oliveira, F.J.R. Paumgartten

Increased EROD activity in the liver of cichlid fishes indicated that Guandu River, the source of drinking water supply for Rio de Janeiro is polluted by CYP1A-inducing chemicals.

239 Determination of trace triazine and chloroacetamide herbicides in tile-fed drainage ditch water using solid-phase microextraction coupled with GC–MS
C. Rocha, E.A. Pappas, C.-h. Huang

Solid-phase microextraction methods were successfully developed to quantify low levels of herbicides in tile-fed drain water by gas chromatography–mass spectrometry.

245 Effect of microbial activity, soil water content and added copper on the temporal distribution patterns of HCB and DDT among different soil organic matter fractions
J.-j. Zhang, B. Wen, X.-q. Shan

Biological activity, soil moisture and Cu have different effects on HCB and DDT transfer from FA, HA and BHA to lipid and IR fractions.

Short communication

253 **Interpreting nitrogen pollution thresholds for sensitive habitats: The importance of concentration versus dose**
I.S.K. Pearce, R. Van der Wal

Nitrate and ammonium concentration in wet deposition detrimentally impacted a sensitive pollution indicator species irrespective of the nitrogen dose.

Corrigendum

257 **Corrigendum to "Introduction" [Environ. Pollut. 148 (2007) 693–694]**
J.A.C. Barth, P. Grathwohl, K.C. Jones

Number 2 2008

Regular papers

259 **Effects of prevalent freshwater chemical contaminants on in vitro growth of *Escherichia coli* and *Klebsiella pneumoniae***
J. Higgins, C. Hohn

Using a microtiter plate assay, E. coli and Klebsiella bacteria were exposed to a panel of common chemical pollutants of fresh water; only ethylene glycol and 2,4-D inhibited bacterial replication.

267 **Comparison of regression coefficient and GIS-based methodologies for regional estimates of forest soil carbon stocks**
J. Elliott Campbell, J.C. Moen, R.A. Ney, J.L. Schnoor

Large differences in estimates of soil organic carbon stocks and annual changes in stocks for Wisconsin forestlands indicate a need for validation from forthcoming forest surveys.

274 **Visible leaf injury in young trees of *Fagus sylvatica* L. and *Quercus robur* L. in relation to ozone uptake and ozone exposure. An Open-Top Chambers experiment in South Alpine environmental conditions**
G. Gerosa, R. Marzuoli, R. Desotgiu, F. Bussotti, A. Ballarin-Denti

Ozone fluxes are more accurate than AOT40 exposure index in predicting ozone visible foliar injury onset on beech seedlings in South Alpine environmental conditions.

285 **Polycyclic aromatic hydrocarbons in ash: Determination of total and leachable concentrations**
A. Enell, F. Fuhrman, L. Lundin, P. Warfvinge, G. Thelin

Total PAH content and the particle weathering rate are more important influences on the rate of migration of PAHs from recycled wood ash into forest soils than leaching.

293 **An assessment of metal contamination along the Irish coast using the seaweed *Ascophyllum nodosum* (Fucales, Phaeophyceae)**
L. Morrison, H.A. Baumann, D.B. Stengel

This paper provides details of an easily applicable, cost-effective and ecologically relevant approach to assessing the degree of metal contamination in coastal environments.

304 **Trends in airborne particulates in Stuttgart, Germany: 1972–2005**
V. Cercasov, V. Wulfmeyer

In Stuttgart, Germany, during 1972–2005 the air pollution shows a clearly decreasing trend, especially for Pb and Br.

314 **Global fallout Pu recorded in lacustrine sediments in Lake Hongfeng, SW China**
J. Zheng, F. Wu, M. Yamada, H. Liao, C. Liu, G. Wan

Fallout Pu recorded in sediments can be used for quick dating of recent sediments to understand the pollution history of environmental contaminants.

322 **Endocrine modulation and toxic effects of two commonly used UV screens on the aquatic invertebrates *Potamopyrgus antipodarum* and *Lumbriculus variegatus***
C. Schmitt, M. Oetken, O. Dittberner, M. Wagner, J. Oehlmann

UV screens can have a significant impact on reproduction and development of aquatic invertebrates.

330 **Use of the modified BCR three-step sequential extraction procedure for the study of trace element dynamics in contaminated soils**
M. Pueyo, J. Mateu, A. Rigol, M. Vidal, J.F. López-Sánchez, G. Rauret

The modified BCR three-step sequential extraction procedure has proved a useful prediction tool for short- and long-term mobility of trace elements in contaminated soils.

342 **Characterisation of esterases as potential biomarkers of pesticide exposure in the lugworm *Arenicola marina* (Annelida: Polychaeta)**
M.L. Hannam, J.A. Hagger, M.B. Jones, T.S. Galloway

This paper provides a preliminary characterisation of esterase enzyme activities in the tissues and body fluids of the sediment dwelling worm Arenicola marina and explores their potential use as biomarkers of organophosphorus pesticide exposure in the marine environment.

351 **Nitrous oxide emission from polyculture constructed wetlands: Effect of plant species**
Y. Wang, R. Inamori, H. Kong, K. Xu, Y. Inamori, T. Kondo, J. Zhang

Zizania latifolia has a large contribution to global warming.

361 ***Tibouchina pulchra* (Cham.) Cogn., a native Atlantic Forest species, as a bio-indicator of ozone: Visible injury**
C.M. Furlan, R.M. Moraes, P. Bulbovas, M.J. Sanz, M. Domingos, A. Salatino

Tibouchina pulchra, a tropical species widely used in Brazilian landscapes as an ornamental tree, is a potential sensitive bio-indicator of ozone air pollution.

366 **Characteristics of organic phosphorus fractions in different trophic sediments of lakes from the middle and lower reaches of Yangtze River region and Southwestern Plateau, China**
R. Zhang, F. Wu, C. Liu, P. Fu, W. Li, L. Wang, H. Liao, J. Guo

Organic phosphorus fractions in sediments from 6 different trophic Chinese lakes were characterized using an improved fractionation scheme.

373 **Contribution of heavy metals and As-loaded lupin root mineralization to the availability of the pollutants in multi-contaminated soils**
S. Vázquez, R.O. Carpena, M.P. Bernal

Maintaining soil pH close to/around 5.5 after lupin culture will be essential for long-time phytostabilization of soils contaminated with heavy metals and arsenic.

380 **Elevated ozone affects the genetic composition of *Plantago lanceolata* L. populations**
R. Kölliker, S. Bassin, D. Schneider, F. Widmer, J. Fuhrer

Five years of exposure indicated a small but significant influence of elevated O_3 on genetic composition and diversity of Plantago lanceolata L. populations.

387 **Temporal variations of surface water quality in urban, suburban and rural areas during rapid urbanization in Shanghai, China**
J. Wang, L. Da, K. Song, B.-L. Li

An integrated pollution index documents the deterioration of water quality in greater Shanghai, recently most serious in rural sections.

394 **Short- and medium-term effects of experimental nitrogen fertilization on arthropods associated with *Calluna vulgaris* heathlands in north-west Spain**
D. Cuesta, A. Taboada, L. Calvo, J.M. Salgado

We observed consistent species-level and variable trophic-group responses to nitrogen addition in one of the southern-most locations for Calluna vulgaris heathlands within Europe.

403 **Seasonal and diurnal gas exchange differences in ozone-sensitive common milkweed (*Asclepias syriaca* L.) in relation to ozone uptake**
C. Bergweiler, W.J. Manning, B.I. Chevone

Temporal variation in physiological processes underlying diurnal and seasonal ozone uptake are described for a key ozone bioindicator species of North America.

416 **Persistent organic pollutants in soils and sediments from James Ross Island, Antarctica**
J. Klánová, N. Matykiewiczová, Z. Máčka, P. Prošek, K. Láska, P. Klán

A survey of soil, sediment and air contamination in James Ross Island, Antarctica serves as a baseline study for evaluation of an anthropogenic impact of the new research facility.

424 **Development of phenanthrene catabolism in natural and artificial soils**
A.H. Rhodes, J. Hofman, K.T. Semple

Natural and artificial soils display different phenanthrene mineralisation profiles suggesting that the extrapolation of results from artificial soils to real/natural soils may not be straightforward.

431 **Persistent organic pollutants (POPs) in Caspian seals of unusual mortality event during 2000 and 2001**
N. Kajiwara, M. Watanabe, S. Wilson, T. Eybatov, I.V. Mitrofanov, D.G. Aubrey, L.S. Khuraskin, N. Miyazaki, S. Tanabe

POPs in seals are high enough to induce immunosuppression.

443 **Seasonal air–water exchange fluxes of polychlorinated biphenyls in the Hudson River Estuary**
S. Yan, L.A. Rodenburg, J. Dachs, S.J. Eisenreich

Investigation of the air–water exchange of PCBs in the Hudson River Estuary suggests that PCBs with 5 or fewer chlorines undergo net volatilization.

452 **Mass fluxes and spatial trends of xenobiotics in the waters of the city of Halle, Germany**
F. Reinstorf, G. Strauch, K. Schirmer, H.-R. Gläser, M. Möder, R. Wennrich, K. Osenbrück, M. Schirmer

Xenobiotics are ubiquitous in the investigated urban aquatic system and are quantified by a large scale mass balance to find spatial trends.

461 **Bulk deposıition of polycyclic aromatic hydrocarbons (PAHs) in an industrial site of Turkey**
F. Esen, S. Sıddık Cindoruk, Y. Tasdemir

Bulk deposition fluxes of PAHs were measured in an industrial area and deposition velocities of PAHs were reported.

468 **Fate and behaviour of phenanthrene in the natural and artificial soils**
J. Hofman, A. Rhodes, K.T. Semple

Artificial soils show substantially different fate and behaviour of phenanthrene than natural soils, which cannot be easily extrapolated or modelled.

476 **Endocrine disruptors in freshwater streams of Hesse, Germany: Changes in concentration levels in the time span from 2003 to 2005**
K. Quednow, W. Püttmann

Concentrations of 4-nonylphenols in rivers of Hessisches Ried in Germany decreased in the sampling period from September 2003 to September 2005.

484 Spatial and temporal variations of mercury levels in Okefenokee invertebrates: Southeast Georgia
B.M. George, D. Batzer

This study measured mercury levels in invertebrates and found the highest levels in amphipods.

491 Influence of fipronil compounds and rice-cultivation land-use intensity on macroinvertebrate communities in streams of southwestern Louisiana, USA
S.V. Mize, S.D. Porter, D.K. Demcheck

Aquatic macroinvertebrate commmunities in southwestern Louisiana streams respond to a gradient of fipronil compounds in water.

Number 3 2008

Regular papers

505 Assessment of sediment contamination in Casco Bay, Maine, USA
T.L. Wade, S.T. Sweet, A.G. Klein

Sediment studies indicate decadal decreases for many chemical contaminants in Casco Bay.

522 Cadmium accumulation and its effect on the in vitro growth of woody fleabane and mycorrhized white birch
R. Fernández, A. Bertrand, A. Casares, R. García, A. González, R.S. Tamés

The high accumulation of Cd observed in selected clones of Dittrichia viscosa and mycorrhized Betula celtiberica grown in vitro implies a potential application for phytoextraction.

530 Response of plankton communities in freshwater pond and stream mesocosms to the herbicide metazachlor
S. Mohr, M. Feibicke, R. Berghahn, R. Schmiediche, R. Schmidt

Metazachlor at levels >5μgL^{-1} is likely to cause long-lasting effects in macrophyte dominated aquatic communities.

543 Accumulation of nine metals and one metalloid in the tropical scallop *Comptopallium radula* from coral reefs in New Caledonia
M. Metian, P. Bustamante, L. Hédouin, M. Warnau

This study investigates metal accumulation behaviour in the tropical scallop Comptopallium radula and preliminary risk assessment for consumers.

553 Levels of dioxins and furans in urban surface soil in Trondheim, Norway
M. Andersson, R.T. Ottesen

PCDD/F levels in soil range between 0.16 and 14 ng I-TEQ kg^{-1} where the main sources are waste incinerators, domestic wood burning and historical sources.

559 Heavy metal accumulation and phytostabilisation potential of tree fine roots in a contaminated soil
I. Brunner, J. Luster, M.S. Günthardt-Goerg, B. Frey

Long-term exposed fine roots of trees are well adapted to soils with high heavy metal contents, but their phytostabilisation capabilities are rather low.

569 Pollution level, phase distribution and health risk of polycyclic aromatic hydrocarbons in indoor air at public places of Hangzhou, China
H. Lu, L. Zhu, S. Chen

Concentrations of PAHs in the air of selected public places in Hangzhou correspond to 10^{-3} life-time lung cancer risk.

576 Effects of pesticides on community composition and activity of sediment microbes – responses at various levels of microbial community organization
A. Widenfalk, S. Bertilsson, I. Sundh, W. Goedkoop

Molecular techniques revealed pesticide-induced changes at lower levels of microbial community organization that were not detected by community-level end points.

585 Arsenate tolerance in *Silene paradoxa* does not rely on phytochelatin-dependent sequestration
M. Arnetoli, R. Vooijs, W. ten Bookum, F. Galardi, C. Gonnelli, R. Gabbrielli, H. Schat, J.A.C. Verkleij

Neither decreased uptake nor phytochelatins seem to play a role in the As tolerance in Silene paradoxa.

592 Determination of Hg and diet identification in otter (*Lontra longicaudis*) feces
C.F. Josef, L.R. Adriano, E.J. De França, G.G. Arantes de Carvalho, J.R. Ferreira

The importance of otter feces preparation for Hg analysis, focusing the food web.

597 The dependence of the β coefficient of REA system with dynamic deadband on atmospheric conditions
T. Grönholm, S. Haapanala, S. Launiainen, J. Rinne, T. Vesala, Ü. Rannik

This paper presents the effect of deadband width and atmospheric stability on the numerical value of empirical β coefficient related to REA system with dynamic deadband.

604 Levels and bioaccumulation of organochlorine pesticides (OCPs) and polybrominated diphenyl ethers (PBDEs) in fishes from the Pearl River estuary and Daya Bay, South China
L. Guo, Y. Qiu, G. Zhang, G.J. Zheng, P.K.S. Lam, X. Li

Bioconcentration was suggested to be responsible for the accumulation of OCPs and PBDEs in the lower trophic organisms of subtropical waters.

612 Effects of nickel chloride and oxygen depletion on behaviour and vitality of zebrafish (*Danio rerio*, Hamilton, 1822) (Pisces, Cypriniformes) embryos and larvae
C. Kienle, H.-R. Köhler, J. Filser, A. Gerhardt

Increasing concentrations of nickel chloride and decreasing concentrations of oxygen lead to reduced vitality and locomotory activity in Danio rerio embryos and larvae.

621 Tissue-specific congener composition of organohalogen and metabolite contaminants in East Greenland polar bears (*Ursus maritimus*)
W.A. Gebbink, C. Sonne, R. Dietz, M. Kirkegaard, F.F. Riget, E.W. Born, D.C.G. Muir, R.J. Letcher

Tissues-specific (adipose tissue, liver, brain and blood) differences exist for the congener patterns of PCBs, PBDEs and their metabolites/degradation products in East Greenland polar bears.

630 Field comparison of disjunct and conventional eddy covariance techniques for trace gas flux measurements
J. Rinne, T. Douffet, Y. Prigent, P. Durand

Results from the first field intercomparison between a new state-of-the-art trace gas flux measurement technique and the direct eddy covariance measurements are reported in this paper.

636 Heavy metal accumulation in *Littoraria scabra* along polluted and pristine mangrove areas of Tanzania
H. De Wolf, R. Rashid

Anthropogenic activities result in heavy metal accumulation and adverse morphological effects in the mangrove gastropod Littoraria scabra.

644 Application and test of a simple tool for operational footprint evaluations
A. Neftel, C. Spirig, C. Ammann

A simple tool for operational footprint calculations is presented and its reliability is assessed using CO_2 flux measurements in a patchy agricultural landscape.

653 Flux and concentration footprint modelling: State of the art
T. Vesala, N. Kljun, Ü. Rannik, J. Rinne, A. Sogachev, T. Markkanen, K. Sabelfeld, Th. Foken, M.Y. Leclerc

The estimation of footprints for the real world with a multitude of interesting gaseous and particulate substances remains a complex problem.

667 Particle fluxes above forests: Observations, methodological considerations and method comparisons
S.C. Pryor, S.E. Larsen, L.L. Sørensen, R.J. Barthelmie

Number fluxes of ultra-fine particles over a forest computed using four micro-meteorological techniques are highly correlated but vary in magnitude.

679 Assessment of sediment concentration and nutrient loads in effluents drained from extensively managed fishponds in France
D. Banas, G. Masson, L. Leglize, P. Usseglio-Polatera, C.E. Boyd

Effluent phosphorus loads were higher than those reported for more intensive aquaculture ponds, but the pollutant source was catchments rather than feeds and fertilizers.

686 Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China
S. Khan, Q. Cao, Y.M. Zheng, Y.Z. Huang, Y.G. Zhu

Long-term wastewater irrigation leads to buildup of heavy metals in soils and food crops.

693 Kinetic extractions to assess mobilization of Zn, Pb, Cu, and Cd in a metal-contaminated soil: EDTA *vs.* citrate
J. Labanowski, F. Monna, A. Bermond, P. Cambier, C. Fernandez, I. Lamy, F. van Oort

Kinetically defined metal fractions mimic mobility aspects of heavy metals.

702 Long-term effects of aided phytostabilisation of trace elements on microbial biomass and activity, enzyme activities, and composition of microbial community in the Jales contaminated mine spoils
G. Renella, L. Landi, J. Ascher, M.T. Ceccherini, G. Pietramellara, M. Mench, P. Nannipieri

Organo-mineral amendment and revegetation of a gold mine spoil increased microbial activity but did not increase microbial species richness.

713 A Cu tolerant population of the earthworm *Dendrodrilus rubidus* (Savigny, 1862) at Coniston Copper Mines, Cumbria, UK
R.E. Arnold, M.E. Hodson, C.J. Langdon

Dendrodrilus rubidus at an abandoned Cu mine tolerates high Cu concentrations by accumulating more Cu before toxic effects occur compared to non-native populations.

723 The nitrate leached below maize root zone is available for deep-rooted wheat in winter wheat–summer maize rotation in the North China Plain
S.-L. Zhou, Y.-C. Wu, Z.-M. Wang, L.-Q. Lu, R.-Z. Wang

Deep-rooted wheat can recycle nitrate leached from maize root zone in winter wheat–summer maize rotation system.

731 Susceptibility to acidic precipitation contributes to the decline of the terricolous lichens *Cetraria aculeata* and *Cetraria islandica* in central Europe
M. Hauck

Artificial acidic precipitation with aqueous sulphur dioxide at pH 2.8–3.5 affects terricolous Cetraria species.

736 Kinetic and dynamic aspects of soil–plant–snail transfer of cadmium in the field
F. Gimbert, M. Mench, M. Coeurdassier, P.-M. Badot, A. de Vaufleury

Toxicokinetics and uptake fluxes can be used to describe the environment contamination by Cd, its bioavailability and transfer to Helix aspersa snails in the field.

Short communication

746 Inorganic arsenic levels in baby rice are of concern
A.A. Meharg, G. Sun, P.N. Williams, E. Adomako, C. Deacon, Y.-G. Zhu, J. Feldmann, A. Raab

Median consumption of organic arsenic levels for UK babies from baby rice is above threshold considered safe.

I Contents of Volume 152